

Name:

Bamfield Number:

Student Number:

Science One Physics Midterm #4

March 18, 2014

Questions 1-8: Multiple Choice: 2 point each

Questions 9-11: Explain your work: 18 points total


Multiple choice answers:

#1	
#2	
#3	
#4	
#5	
#6	
#7	
#8	

Formula sheet at the back (you can remove it)


Question 1: In the picture below, the moving charge will be deflected

- A) into the page.
- B) out of the page
- C) upwards
- D) downwards
- E) None of the above.



Question 2: A magnet sits in a non-uniform magnetic field as shown to the right. Ignoring gravity, we can say that the magnet will

- A) accelerate upward
- B) accelerate downward
- C) experience no net force




Question 3: The picture above shows the actual path that a proton which starts from rest will follow in the presence of certain uniform electric and magnetic fields. We can say that:

- A) the electric field points to the right and the magnetic field points into the page
- B) the electric field points down and the magnetic field points into the page
- C) the electric field points to the right and the magnetic field points downward
- D) the electric field points down and the magnetic field points to the right
- E) the electric field points up and the magnetic field points downward
- F) the electric field points up and the magnetic field points into the page

Question 4: What happens when the switch is closed?


- A) B and C both get brighter
- B) B gets brighter and C gets dimmer
- C) Nothing happens
- D) B gets dimmer and C get brighter
- E) B and C both get dimmer



Question 5: Current flows through a wire that is thick at one end and thin at another end. Which of the following change from the thick end to the thin end?

- A) Current density
- B) Conductivity
- C) Electric field
- D) both A) and B)
- E) both A) and C)
- F) both B) and C)
- G) all of A), B), and C)


Question 6: The picture to the right shows a positive charge sitting in a magnetic field produced by a solenoid. If the current in the solenoid is decreased, we can say that the charge will initially move



- A) to the right
- B) to the left
- C) upward
- D) downward
- E) into the page
- F) out of the page
- G) None of the above: it will stay still.

Question 7: A bunch of Science One would like to increase the EMF of a generator they have built (that spins a coil of wire in a magnetic field). Which of the following will **not** help:

- A) Increasing the magnetic field strength
- B) Increasing the area of the loop
- C) Increasing the number of turns of wire
- D) Decreasing the resistance of the wire



Question 8: When the switch is closed, what best describes what happens in the circuit above?

- A) Light bulbs A and B both suddenly go dim. They then return to their previous brightness.
- B) Light bulbs A and B both suddenly go dim. They slowly return to different brightness than when the switch was closed.
- C) Light bulb B goes out and A suddenly gets brighter. They then slowly return to their previous brightness.
- D) Light bulb B goes out and A suddenly gets brighter. They slowly return to different brightness than when the switch was closed.
- E) Light bulbs A and B both suddenly go brighter. They then return to their previous brightness.
- F) Light bulbs A and B both suddenly go dimmer. They then return to their previous brightness.


Question 10: Give a brief explanation (1-2 sentences) for each of the following:

a) If electrons accelerate in an electric field, why doesn't current increase steadily with time when we apply a voltage to a conductor? **(2 points)**


b) Why does a magnet stick to a refrigerator door if the door isn't a magnet?
(2 points)

c) When we connect a capacitor to a battery, why does current flow out of the battery for a while and then stop? **(2 points)**

Question 11: A charged particle with mass 10^{-15} kg and charge 5×10^{-10} C starts at rest at the bottom of a capacitor in which the electric field is $10,000$ V/m. The particle accelerates upward, passing through a hole in the top plate and entering a uniform magnetic field. If we want the particle to end up in a collector that is 2 cm from the hole, as shown, how strong a magnetic field do we need? (6 points)



Question 12: For reasons unknown, Sally would like to hang up a pair of pants using electromagnetic induction instead of an ordinary clothesline. She builds a conducting frame with a conducting bar that touches the two sides but can move up and down freely without friction. She hangs a damp pair of pants on the bar, and then turns on an increasing uniform magnetic field.



- a) Explain why this allows Sally to hold up the pants.
(3 points)
- b) If the mass of the bar + pants is 1 kg, and the resistance of the loop created by the bar and frame is 0.1 Ohms, determine $B(t)$ so that the pants will stay still (or at least find an equation that $B(t)$ must satisfy). *Ignore any magnetic fields other than the one Sally turns on.*
(3+ points)

FORMULA SHEET

$$a = dv/dt \quad v = dx/dt$$

$$\mathbf{F} = m\mathbf{a} \quad a = v^2/R$$
$$F_r = -dU/dr \quad W = -\Delta U = -\int \mathbf{F} \cdot d\mathbf{r}$$

$$\mathbf{F} = q\mathbf{E} \quad U = qV$$
$$E_r = -dV/dr \quad \Delta V = -\int \mathbf{E} \cdot d\mathbf{r}$$

$$E = kq/r^2 \quad E = \eta/(2\epsilon_0) \quad E = 2kp/r^3 \quad p = qs$$
$$\text{Flux} = Q_{\text{enc}}/\epsilon_0$$

$$k = 9 \times 10^9 \text{ N m}^2/\text{C}^2 \quad \epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/(\text{N m}^2)$$

$$\mathbf{F} = q\mathbf{E} + q \mathbf{v} \times \mathbf{B} \quad \mathbf{F} = I \mathbf{l} \times \mathbf{B} \quad \boldsymbol{\tau} = \boldsymbol{\mu} \times \mathbf{B} \quad \boldsymbol{\mu} = I \mathbf{A}$$

$$\mathbf{B} = \mu_0/(4\pi)q \mathbf{v} \times \mathbf{r} / r^3 \quad \mathbf{B} = \mu_0/(4\pi)I d\mathbf{s} \times \mathbf{r} / r^3 \quad B = (\mu_0 / 2 \pi) I/d \quad B = \mu_0 (N/L) I$$

$$V = IR$$

$$C = Q/V$$

$$P = IV$$

$$R = \rho LA$$

$$\sigma = n_e e^2 \tau / m = 1/\rho$$

$$v_d = e \tau E / m$$

$$I = e n_e A v_d$$

$$Q(t) = Q_0 \exp(-t/RC)$$

$$\epsilon = |d\Phi_m/dt|$$

$$\Phi = \mathbf{B} \cdot \mathbf{A} = BA \cos(\theta)$$

$$\oint \vec{E} \cdot d\vec{s} = -d\Phi_m/dt$$